

Third potential evidence for the new area spectrum based on the Einstein-Kaufman pseudo tensor, a conjecture

August 1, 2011

Brian Kong ¹, Youngsub Yoon ²

¹*Milton Academy, Milton, MA 02186, USA*

²*Harvard University, Cambridge, MA 02138, USA*

Abstract

In our earlier paper “Corrections to the Bekenstein-Hawking entropy and the Hawking radiation spectrum,” arXiv:0910.2755, we provided two concrete numerical evidences for the new area spectrum based on the Einstein-Kaufman pseudo tensor as opposed to the Ashtekar variables: namely, the reproduction of the Bekenstein-Hawking entropy without fixing Immirzi parameter and the reproduction of the Hawking radiation spectrum. In this article, we provide another potential, albeit not concrete, numerical evidence for this new area spectrum; there was a constant in our earlier article, which was inversely proportional to the density of state, and which we could not fix a priori. Nevertheless, in our earlier article, we obtained this constant to be around 172~173 by fitting it to the Planck radiation spectrum. In this article, we propose a mathematical formula that reproduces this value. According to this formula, this constant is around 172.8. Nevertheless, we failed to concretely derive this mathematical formula. Therefore, it is a conjecture.

1 Our Argument

In our earlier paper [1] we derived the number of photons emitting from a black hole in terms of A_{BH} , the area of the black hole and A_{spec} , the area spectrum corresponding to the photon radiated. The formula we obtained was following:

$$dn_{photon} \propto \frac{A_{spec}^2 dA_{spec}}{\sqrt{A_{BH}}(e^{A_{spec}/4} - 1)} \quad (1)$$

where we ignored the unimportant proportionality constant. Of course, the area spectrum A_{spec} , and the frequency of photon radiating f is related by the following formula.

$$\frac{hf}{kT} = \frac{A_{spec}}{4} \quad (2)$$

Therefore, we can easily see that the total energy radiated by a black hole is given by the following formula.

$$E_{total} \propto \int_{A_{min}}^{\infty} \frac{A_{spec}^3 dA_{spec}}{A_{BH}(e^{A_{spec}/4} - 1)} \quad (3)$$

where we ignored the unimportant proportionality constant again and where A_{min} is the lowest area spectrum. Also, in our earlier article we argued that the Hawking radiation spectrum is reproduced if the following formula holds which we numerically showed was the case.

$$dN(A) = \frac{1}{C} \sqrt{A} (e^{A/4} - 1) dA \quad (4)$$

where $N(A)$ is the number of degeneracy for the area A . In our earlier paper [1], we also mentioned that we were not able to derive C in the above equation; all we could do was obtaining C by fitting. We obtained C to be around 172~173. In this paper, we will suggest a formula that reproduces C .

Motivated by (3) we considered the following integration.

$$\int_{A_{min}}^{\infty} \frac{A_{spec}^3 dA_{spec}}{e^{A_{spec}/4} - 1} = G \quad (5)$$

Plugging in for A_{min} for $4\pi\sqrt{2}$, we obtained G to be approximately 542.9. Surprisingly, this value is πC . In other words,

$$C = \frac{1}{\pi} \int_{A_{min}}^{\infty} \frac{A_{spec}^3 dA_{spec}}{e^{A_{spec}/4} - 1} \quad (6)$$

The right-hand side is approximately 172.8 which agrees with the left-hand side C which is around 172~173 as we showed in our earlier paper.

2 Discussion and Conclusions

In this paper, we provided the third potential evidence for our new area spectrum. Even though this evidence is only potential and not concrete as our previous two evidences which we provided in our earlier paper, it would give another undeniable support for our new area spectrum if our formula (6) is concretely derived. We hope that this formula will soon be derived by future research.

References

- [1] B. Kong and Y. Yoon, “Corrections to the Bekenstein-Hawking entropy and the Hawking radiation spectrum,” arXiv:0910.2755 [gr-qc].